

- Hurst, G.D.D., Sharpe, R.G., Broomfield, A.H., Walker, L.E., Majerus, T.M.O., Zakharov, I.A. & Majerus, M.E.N., 1995. Sexually transmitted disease in a promiscuous insect, *Adalia bipunctata* L.. *Ecological Entomology* **20**: 230-236.
- Iperti, G., 1964. Les parasites des Coccinelles aphidiphages dans les Alpes-Maritimes et les Basses-Alpes. *Entomophaga* **9**: 153-180.
- Kuznetsov, V.N., 1993. *Coccinellids (Coleoptera: Coccinellidae) of the Russian Far East*. Dalnauka: Vladivostok.
- Majerus, M.E.N., 1989. *The Cambridge Ladybird Survey – Results Supplement 1984-1989*. Department of Genetics: Cambridge.
- , 1994. *Ladybirds*. New Naturalist Library 81. Harper Collins: London.
- , 1997. Parasitisation of British coccinellids by *Dinocampus coccinellae* (Hymenoptera: Braconidae). *Br. J. Ent. Nat. Hist.* **10**: 15-24.
- Majerus, M.E.N. & Majerus, T.M.O., 1997. Predation of ladybirds by birds in the wild. *Ent. Mon. Mag.* **133**: 55 - 61.
- Marples, N.M., Brakefield, P.M. & Cowie, R.J., 1989. Differences between the 7-spot and 2-spot ladybird beetles (Coccinellidae) in their toxic effects on a bird predator. *Ecological Entomology* **14**: 79-84.
- Morton Jones, F., 1932. Insect coloration and the relative acceptability of insects to birds. *Trans. Ent. Soc. Lond.* **80**: 345-385.
- Pasteels, J.M., Deroe, C., Tursch, B., Brakeman, J.C., Daloze, D. & Hootele, C., 1973. Distribution et activités des alcaloïdes défensifs des Coccinellidae. *J. Insect Physiology* **19**: 1771-1784.
- Pocock, R.I., 1911. On the palatability of some British insects, with notes on the significance of mimetic resemblances. With notes on the experiments by E.B. Poulton. *Proc. Zool. Soc. Lond.* **1911**: 809-868.
- Tursch, B., Daloze, D., Dupont, M., Hootele, C., Kaisin, M., Pasteels, J.M. & Zimmermann, D., 1971. Coccinelline, the defensive alkaloid of the beetle *Coccinella septempunctata*. *Chemia* **25**: 307.
- 
- 

### **A second Scottish record of *Ypsolopha sequella* (Clerck) (Lep.: Yponomeutidae)**

The distribution of *Ypsolopha sequella* extends throughout England and Wales as far as the Scottish border (Agassiz, 1996. in Emmet (Ed.) *The moths and butterflies of Great Britain and Ireland*. **3**:97). A single record by Andrew Buckham from Denholm, Roxburghshire (VC 80) on 26 August 1975 was, until now, the only known incursion of this species across the border. On 16 September 1997 a second Scottish specimen of *Y. sequella* was found at rest on the trunk of a sycamore *Acer pseudoplatanus* tree near the Eastern General Hospital in Leith, Midlothian (VC 83; OS grid ref. NT 2875). It will be interesting to see if further expansion of its range occurs.— K.P. BLAND, 35 Charterhall Road, Edinburgh EH9 3HS and D.M. ROBERTSON, 3 Claremont Park, Leith, Edinburgh EH6 7PH.